15 15

ţħ

, <u>T</u>

ĻΠ

ŧ.

<u>}</u> =

20 mm mm 20

25

30

WHAT is CLAIMED is:

- A method for decreasing serum cholesterol and was Increasing serum HDL in a patient comprising administering to the digestive tract of said patient an effective amount 5 of a composition comprising a viable lactic acid-producing bacteria and a therapeutic agent selected from the group consisting of an\effective amount of a cholesterol-reducing agent and a bifidogenic oligosaccharide.
 - The method of claim 1 wherein said lactic acidproducing bacteria is selected from the group consisting of 10 non-pathogenic members of the Bacillus genus, Lactobacillus, Sporolactobacithus and Bifidobacterium.
 - The method of claim 2 wherein said lactic acidproducing Baci Nus is selected from the group consisting of Bacillus coagulans, Bacillus coagulans Hammer, Bacillus brevis subspecies cagulans and Bacillus laevolacticus.
 - The method of claim 1 wherein said lactic acid-4. producing bacteria is Bacillus coagulans subspecies Hammer.
 - The\method of claim 2 wherein said lactic acidproducing Lact&bacillus is selected from the group consisting of Lactobacillus acidophilus, Lactobacillus casei, Lactobacillus DDS-1, Lactobacillus GG, Lactobacillus rhamnosus, Lactobacillus plantarum, Lactobacillus salivarius and Lactobacillus sporogenes (aka B. coagulans).
 - The metho $A \circ A$ of claim 2 wherein said lactic acidproducing Sporolactobacillus P44.
 - The method of claim 2 wherein said lactic acidproducing Bifidobacteritm is selected from the group consisting of Bifidobacterium adolescentis, Bifidobacterium animalis, Bifidobacterium\bifidum, Bifidobacterium bifidus, Bifidobacterium breve, Bifidobacterium infantis, Bifidobacterium infantus and Bifidobacterium longum.
 - 8. The method of claim 1 wherein said composition

contains 10^5 to 10^{16} viable bacterium per gram of composition.

5

10

1

[] ___15

14

ļ. <u>.</u>

- 9. The method of claim 1 wherein said administering comprises oral ingestion of said composition.
- 10. The method of claim 1 wherein said administering comprises introducing into the digestive tract from 0.1 to 5 grams per day of said composition.
- 11. The method of claim 1 wherein said administering comprises introducing into the digestive tract from 10^8 to 10^{10} viable bacterium per day.

12. The method of claim 11 wherein said administering comprises introducing into the digestive tract from 5×10^8 to 10^9 viable bacterium per day.

- 13. The method of claim 1 wherein said bifidogenic oligosaccharide is selected from the group consisting of fructo-oligosaccharide, gluco-oligosaccharide, and trisaccharide raffinose.
- 14. The method of claim 13 wherein fructooligosaccharide comprises polymers of fructose and glucose having a polymer chain length of from about 4 to 100 sugar units.
- 15. The method of claim 1 wherein said composition comprises about 10 milligrams to about 1 gram of bifidogenic oligosaccharide per gram of composition.
- 25 16. The method of claim 1 wherein said composition comprises from 100 to 500 milligrams of bifidogenic oligosaccharide per gram of composition.
- 17. The method of claim 1 wherein said administering comprises introducing into the digestive tract from 10 milligrams to 20 grams of bifidogenic oligosaccharide per day.
 - 18. The method of claim 17 wherein said administering comprises introducing into the digestive tract from 150

milligrams to 5 grams of bifidogenic oligosaccharide per day.

- 19. The method of claim 1 wherein said cholesterol-reducing agent is selected from the group consisting of a statin, a bile sequestering compound, a fiber product capable of binding cholesterol, niacin and aspirin.
- 20. The method of claim 19 wherein said statin is selected from the group consisting of cerivastatin, fluvastatin, lovastatin, pravastatin and simvastatin.
- 10 21. The method of claim 20 wherein said administering comprises introducing into the digestive tract from 10 to 80 milligrams of statin per day.
 - 22. The method of claim 19 wherein said bile sequestering compound is selected from the group consisting of colestipol and cholestyramine.

١...

ξĦ

[]

25

....... 15

- 23. The method of claim 22 wherein said administering comprises introducing into the digestive tract from 1 to 20 grams of bile sequestering compound per day.
- 24. The method of claim 19 wherein said fibrin is selected from the group consisting of gemfibrozil, fenofibrate, psyllium, bran, glucomannan and Jerusalem artichoke flour.
- 25. The method of claim 24 wherein said administering comprises introducing into the digestive tract from 500 milligrams to 50 grams of fibrin per day.
- 26. The method of claim 1 wherein said compost further comprises a cholic acid complexation agent.
 - 27. The method of claim 26 wherein said complexation agent is selected from the group consisting of a metal salt of calcium, chromium, copper, iodine, iron, magnesium, manganese, potassium, sodium and zinc.
 - 28. The method of claim 27 wherein said metal salt is provided in the form of calcium citrate, potassium

gluconate, magnesium citrate or chromium picollinate.

- 29. The method of claim 1 wherein said composition further comprises a food substance, flavoring, vitamin or mineral.
- 30. The method of claim 1 wherein said patient is at risk for atherosclerosis, arterial sclerosis, myocardial infarction, heart attack, diabetes, coronary heart disease, angina pectoris or unstable angina.
- 10 31. A therapeutic composition for reduction of serum cholesterol comprising a viable lactic acid-producing bacteria and a therapeutic agent selected from the group consisting of an effective amount of a cholesterol-reducing agent and a bifidogenic oligosaccharide.

Annual Carlo and the second and the second

THE THE REPORT

ij

30

15

- 32. The composition of claim 31 wherein said lactic acid-producing bacteria is selected from the group consisting of non-pathogenic members of the Bacillus genus, Lactobacillus, Sporolactobacillus and Bifidobacterium.
- 33. The composition of claim 32 wherein said lactic acid-producing Bacillus is selected from the group consisting of Bacillus coagulans, Bacillus coagulans Hammer, Bacillus brevis subspecies coagulans and Bacillus laevolacticus.
- 34. The composition of claim 31 wherein said lactic 25 acid-producing bacteria is Bacillus coagulans subspecies Hammer.
 - 35. The composition of claim 32 wherein said lactic acid-producing Lactobacillus is selected from the group consisting of Lactobacillus acidophilus, Lactobacillus casei, Lactobacillus DDS-1, Lactobacillus GG, Lactobacillus rhamnosus, Lactobacillus plantarum, Lactobacillus salivarius and Lactobacillus sporogenes (aka B. coagulans).
 - 36. The composition of claim 32 wherein said lactic

acid-producing Sporolactobacillus is Sporolactobacillus P44.

- 37. The composition of claim 32 wherein said lactic acid-producing Bifidobacterium is selected from the group consisting of Bifidobacterium adolescentis, Bifidobacterium animalis, Bifidobacterium bifidum, Bifidobacterium bifidus, Bifidobacterium breve, Bifidobacterium infantis, Bifidobacterium infantus and Bifidobacterium longum.
 - 38. The composition of claim 31 wherein said composition contains 10^5 to 10^{10} viable bacterium per gram of composition.

10

The second of th

ţñ

17.4

<u>(</u>520

- 39. The composition of claim 31 wherein said bifidogenic oligosaccharide is selected from the group consisting of fructo-oligosaccharide, gluco-oligosaccharide, and trisaccharide raffinose.
- 40. The composition of claim 39 wherein fructooligosaccharide comprises polymers of fructose and glucose having a polymer chain length of from about 4 to 100 sugar units.
- 41. The composition of claim 31 wherein said composition comprises about 10 milligrams to about 1 gram of bifidogenic oligosaccharide per gram of composition.
- 42. The composition of claim 31 wherein said composition comprises from 100 to 500 milligrams of bifidogenic oligosaccharide per gram of composition.
- 25 43. The composition of claim 31 wherein said cholesterol-reducing agent is selected from the group consisting of a statin, a bile sequestering compound, a fiber product capable of binding cholesterol, niacin and aspirin.
- 30 44. The composition of claim 43 wherein said statin is selected from the group consisting of cerivastatin, fluvastatin, lovastatin, pravastatin and simvastatin.
 - 45. The composition of claim 43 wherein said

composition comprises from 1 to 80 milligrams of statin per gram of composition.

- 46. The composition of claim 43 wherein said bile sequestering compound is selected from the group consisting of colestipol and cholestyramine.
 - 47. The composition of claim 43 wherein said composition comprises from 0.1 to 0.8 grams of bile sequestering compound per gram of composition.

5

Green, gare, green, steel of green, steel and Mark Mark Mark

15

20 m state in the street s

}- ±

- 48. The composition of claim 43 wherein said fibrin is selected from the group consisting of gemfibrozil, fenofibrate, psyllium, bran, glucomannan and Jerusalem artichoke flour.
 - 49. The composition of claim 43 wherein said composition comprises from 10 milligrams to 0.5 grams of fibrin per gram of composition.
 - 50. The composition of claim 31 wherein said compost further comprises a cholic acid complexation agent.
 - 51. The composition of claim 50 wherein said complexation agent is selected from the group consisting of a metal salt of calcium, chromium, copper, iodine, iron, magnesium, manganese, potassium, sodium and zinc.
 - 52. The composition of claim 51 wherein said metal salt is provided in the form of calcium citrate, potassium gluconate, magnesium citrate or chromium picollinate.
- 25 53. The composition of claim 31 wherein said composition further comprises a food substance, flavoring, vitamin or mineral.
- 54. A therapeutic system for reducing serum cholesterol comprising a container comprising a label and a composition comprising a culture of viable lactic acid-producing bacteria and a therapeutic agent selected from the group consisting of an effective amount of a cholesterol-reducing

agent and a bifidogenic oligosaccharide, wherein said label comprises instructions for use of the composition for reduction of serum cholesterol.

- 55. The system of claim 54 wherein said lactic acid-producing bacteria is selected from the group consisting of non-pathogenic members of the Bacillus genus, Lactobacillus, Sporolactobacillus and Bifidobacterium.
- 56. The system of claim 55 wherein said lactic acidproducing Bacillus is selected from the group consisting of
 Bacillus coagulans, Bacillus coagulans Hammer, Bacillus
 brevis subspecies coagulans and Bacillus laevolacticus.
 - 57. The system of claim 55 wherein said lactic acidproducing bacteria is Bacillus coagulans subspecies Hammer.
 - 58. The system of claim 55 wherein said lactic acid-producing Lactobacillus is selected from the group consisting of Lactobacillus acidophilus, Lactobacillus casei, Lactobacillus DDS-1, Lactobacillus GG, Lactobacillus rhamnosus, Lactobacillus plantarum, Lactobacillus salivarius and Lactobacillus sporogenes (aka B. coagulans).

h ±

25

30

- 59. The system of claim 55 wherein said lactic acid-producing Sporolactobacillus is Sporolactobacillus P44.
- 60. The system of claim 55 wherein said lactic acidproducing Bifidobacterium is selected from the group
 consisting of Bifidobacterium adolescentis, Bifidobacterium
 animalis, Bifidobacterium bifidum, Bifidobacterium bifidus,
 Bifidobacterium breve, Bifidobacterium infantis,
 Bifidobacterium infantus and Bifidobacterium longum.
- 61. The system of claim 54 wherein said composition contains 10^5 to 10^{10} viable bacterium per gram of composition.
- 62. The system of claim 54 wherein said bifidogenic oligosaccharide is selected from the group consisting of fructo-oligosaccharide, gluco-oligosaccharide, and

trisaccharide raffinose.

5

THE REAL RESIDENCE OF THE STATE OF THE STATE

13

14

15

20

- 63. The system of claim 62 wherein fructooligosaccharide comprises polymers of fructose and glucose having a polymer chain length of from about 4 to 100 sugar units.
- 64. The system of claim 54 wherein said composition comprises about 10 milligrams to about 1 gram of bifidogenic oligosaccharide per gram of composition.
- 65. The system of claim 54 wherein said composition comprises from 100 to 500 milligrams of bifidogenic oligosaccharide per gram of composition.
 - 66. The system of claim 54 wherein said cholesterol-reducing agent is selected from the group consisting of a statin, a bile sequestering compound, a fiber product capable of binding cholesterol, niacin and aspirin.
 - 67. The system of claim 66 wherein said statin is selected from the group consisting of cerivastatin, fluvastatin, lovastatin, pravastatin and simvastatin.
 - 68. The system of claim 66 wherein said composition comprises from 1 to 80 milligrams of statin per gram of composition.
 - 69. The system of claim 66 wherein said bile sequestering compound is selected from the group consisting of colestipol and cholestyramine.
- 70. The system of claim 66 wherein said composition comprises from 0.1 to 0.8 grams of bile sequestering compound per gram of composition.
 - 71. The system of claim 66 wherein said fibrin is selected from the group consisting of gemfibrozil,
- 30 fenofibrate, psyllium, bran, glucomannan and Jerusalem artichoke flour.
 - 72. The system of claim 66 wherein said composition comprises from 10 milligrams to 0.5 grams of fibrin per gram

of composition.

- The system of claim 54 wherein said compost further comprises a cholic acid complexation agent.
- 74. The system of claim 54 wherein said complexation agent is selected from the group consisting of a metal salt of calcium, chromium, copper, iodine, iron, magnesium, manganese, potassium, sodium and zinc.
 - The system of claim 74 wherein said metal salt is provided in the form of calcium citrate, potassium gluconate, magnesium citrate or chromium picollinate.
 - The system of claim 54 wherein said composition further comprises a food substance, flavoring, vitamin or mineral.

5

then and and a great of after the form of the first of th